

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



29-0-0 W/V
Guaranteed Analysis

ANALYSIS

Nitrogen (N)

19.43% Urea Nitrogen

9.57% Other Water Soluble Nitrogen*

Derived from: Urea-Triazone Solution

*9.57% Slowly Available Nitrogen from Urea-Triazone Solution

W/V%
29.00

GENERAL INFORMATION

Maximum N-Pact™ is a foliar Fertiliser that contains Triazone, a patented slow-release nitrogen compound. **Maximum N-Pact** may be applied whether undiluted or diluted (with water) for aerial or ground application. Use sufficient carrier to adequately cover crop foliage.

Phytotoxicity: Plant and leaf injury may occur on some crops when certain weather and growing conditions are present. User assumes all risks of use and handling.

DIRECTIONS FOR USE SUGGESTED APPLICATION RATES

CROP	SUGGESTED APPLICATION RATES AND TIMING
Almonds	10 to 20 L/ha at full leaf. Repeat at early nut expansion.
Apples	10 to 15 L/ha Apply in a minimum of 500 litres of water spray solution. Begin at first full leaf and apply as needed during the growing season.
Apricots	10 to 15 L/ha prior to fruit set.
Asparagus	10 to 20 L/ha beginning at mid-fern development, then at 14 to 21 day intervals.
Beans (Dry)	Single Application: 20 L/ha at early pod formation. Multiple Applications: 10 to 15 L/ha at early flower and repeat in 10 to 14 days.
Beans (green, lima)	10 to 15 L/ha at early flower and repeat in 7 to 10 days.
Blueberries	10 to 20 L/ha at early fruit set and repeat at early fruit color.
Broccoli	10 to 20 L/ha. First application after thinning, then 3 weeks before head formation and repeat in 7 to 10 days.
Cabbage	10 to 20 L/ha. First application after thinning, then at head formation and repeat after 14 to 21 days.
Caneberries	10 to 15 L/ha prior to fruit set.
Canola	10 to 20 L/ha at pre-bloom.
Carrots	10 to 15 L/ha. First application when plants are 7.5 to 15 cm tall, then repeat at 21 day intervals or as required.
Cauliflower	10 to 20 L/ha. First application after thinning or transplant, other applications at early head set and repeat at 10 to 14 day intervals.
Celery	10 to 15 L/ha. First application when plants are 20 to 30 cm tall and repeat at 10 to 14 day intervals.
Cherries	10 to 20 L/ha prior to fruit set.
Christmas Trees,	10 to 20 L/ha when sufficient foliage is present. Repeat as needed.
Ornamental and Nursery Stock	
Citrus	10 to 20 L/ha at early bloom and repeat after fruit set.
Corn	10 to 20 L/ha before detasseling and repeat after pollination.
Corn (Sweet)	10 to 15 L/ha. First application when plants are 30 to 60 cm high, then at tassel emergence and repeat after pollination.
Cotton (After Seeding Stage)	5 to 10 L/ha may be applied in combination with crop protection chemicals and growth regulators.
Cotton (Boll Development)	5 to 10 L/ha at early boll formation and repeat at 14 to 21 day intervals.
Cotton (Seeding)	5 to 10 L/ha after first true leaves appear.
Cranberries	10 to 15 L/ha at hook stage and repeat after fruit set.

Cucumbers	Single Application: 20 to 40 L/ha at early fruit set. Multiple Applications: 10 to 15 L/ha at early flower and repeat at 10 to 14 day intervals.
Eggplant	10 to 15 L/ha at bud stage and repeat at 10 to 14 day intervals.
Flax	10 to 20 L/ha at early boll development.
Grapes (Raisin)	5 to 10 L/ha when sufficient foliage is present. Repeat as needed.
Grapes (Table)	5 to 10 L/ha prior to fruit set.
Grapes (Wine)	5 to 10 L/ha when sufficient foliage is present. Repeat as needed.
Grass (Seed Production)	20 to 40 L/ha at seed head elongation.
Hazelnuts	Single Application: 20 to 40 L/ha at early nut filling. Multiple Applications: 10 to 15 L/ha at early leaf expansion and repeat at 14 to 21 day intervals.
Hops	10 to 15 L/ha before cone development.
Lentils	10 to 15 L/ha at early flowering and repeat in 10 to 14 day intervals.
Lettuce	10 to 15 L/ha after thinning, then at early head formation and repeat at 10 to 14 day intervals.
Lucerne	10 to 15 L/ha Apply after each cutting when sufficient foliage is present.
Nectarines	10 to 20 L/ha prior to fruit set.
Okra	10 to 15 L/ha at bud stage and repeat at 10 to 14 day intervals.
Olives	10 to 15 L/ha at early fruit development and repeat as needed.
Onions	10 to 15 L/ha at mid-set development and repeat at 14 to 21 day intervals.
Other Crops	5 to 40 L/ha when sufficient foliage is present. Try on a small area until more experience and trials have been completed to determine if higher rate is desirable.
Pasture grazing & fodder crops	10L/ha from early seeding stage when sufficient foliage is present. With further applications as required.
Pastures	10L/ha from early seeding stage. With further applications as required.
Peaches	10 to 20 L/ha prior to fruit set.
Peanuts	Single Application: 20 to 40 L/ha at mid-tuber development. Multiple Applications: 10 to 15 L/ha at early bloom and repeat at 14 to 21 day intervals until pods are filled.
Pears	10 to 15 L/ha prior to fruit set or post-harvest.
Peas	10 to 15 L/ha at early flowering and repeat in 10 to 14 days.
Pecans	10 to 15 L/ha at full leaf. Repeat at early nut expansion.
Peppers	10 to 15 L/ha at early fruit set and repeat at 10 to 14 day intervals.
Plums	10 to 20 L/ha prior to fruit set.
Potatoes	10 to 15 L/ha 4-6 leaf stage. With further applications as required.
Prunes	5 to 10 L/ha at full leaf. Repeat as needed.
Rice	10 to 20 L/ha at panicle initiation. Repeat as required.
Rock Melon	10 to 20 L/ha at early flowering and repeat in 7 to 10 days.
Sorghum (Forage and Grain)	10 to 15 L/ha after pollination.
Soybeans	10 to 15 L/ha at early flower and repeat in 14 to 21 days.
Spinach	10 to 20 L/ha when sufficient foliage is present and repeat at 14 to 21 days.
Squash	10 to 20 L/ha at early fruit set and repeat at 10 to 14 day intervals.
Strawberries	5 to 10 L/ha at early flowering and repeat every 14 days through harvest. Initiate fall application when new growth reaches 7.5 cm in height.
Sugar Beets	20 L/ha at 10 to 12 leaf stage and repeat at 20 leaf stage.
Sunflower	10 to 15 L/ha when outer seeds start to fill, repeat in 10 to 14 days.
Sweet Potatoes	10 to 15 L/ha 4-6 leaf stage. With further applications as required.
Tomatoes (Fresh)	Single Application: 20 L/ha 10 to 14 days after full bloom. Multiple Applications: 10 to 15 L/ha at full bloom and repeat at 10 to 14 day intervals until harvest is near completion.
Tomatoes (Process)	Single Application: 20 L/ha 10 to 14 days after full bloom. Multiple Applications: 10 to 15 L/ha at full bloom and repeat at 10 to 14 day intervals until 2 weeks before harvest.
Watermelons	10 to 20 L/ha at early flowering and repeat in 7 to 10 days.
Wheat or Barley	10 to 20 L/ha at tiller to seed head formation.

Maximum N-Pact has been applied undiluted to some types of ornamental and nursery stock. Since these plants vary widely in growth habit and leaf texture, some types are more susceptible to leaf or tip burn than others. It is recommended that the user try only a few plants at undiluted or concentrated rates and observe for a week before general applications are made.

MIXING INSTRUCTIONS

Maximum N-Pact mixes easily with water, as follows:

1. Fill tank half-full with water and start agitation.
2. Add recommended amounts of any flowable materials, any emulsifiable materials, any soluble powders and/or water soluble fertilizers. (All soluble powders and/or water soluble fertilizers should be pre-dispersed in water before adding to the spray tank.) then add compatible micronutrients ensure all product is in suspension before adding **Maximum N-Pact**.

STORAGE and DISPOSAL

Storage: Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. DO NOT contaminate water, food, or feed by storage or disposal.

Disposal: DO NOT reuse container. Completely empty container into application equipment.

Thoroughly rinse container, adding rinse water to the spray tank.

Handling: Avoid breathing Fertiliser mists. Avoid prolonged or repeated skin contact. Avoid eye contact. Wash hands thoroughly after handling this product.

FIRST AID

If in Eyes: Flush with clean running water for at least 15 minutes. If irritation persists, get medical attention. **If on Skin:** Remove contaminated clothing. Wash thoroughly with soap and water. If irritation persists, or if rash develops, get medical attention. **If Swallowed:** If conscious, drink 1 or 2 glasses of water. Get medical attention. Never induce vomiting or give anything by mouth to an unconscious person. **If Inhaled:** Remove to fresh air. If irritation persists, get medical attention.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet, which is available from the supplier.

CONDITIONS OF SALE: Any provisions or rights (under the Competition and Consumer Act 2010 and/or relevant state legislation) which cannot be excluded by those statutes or by law are not intended to be excluded by these Conditions of Sale. Subject to the foregoing, all warranties, conditions, rights and remedies (expressed or implied under common law, statute, or otherwise) in relation to the sale, supply, use and/or application of this product are excluded. This product must be used strictly as directed and in accordance with all instructions appearing on the label and other reference material. To the extent allowed by statute, Loveland Products, Inc. and/or its affiliates shall not accept any liability whatsoever (including consequential loss), howsoever arising (including negligence), for any damage, injury or death connected with the sale, supply, use and/or application of this product.

Not classified as Hazardous according to the Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia criteria.

MAXIMUM N-PACT® is a registered trademark of Loveland AgriProducts

Manufactured for: Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113
Tel (02) 9889 5400 • **Product Support Tel: 1800 44 88 92**

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE
FOR SPECIALIST ADVICE IN
AN EMERGENCY DIAL
1800 033 111
24 HOURS AUSTRALIA WIDE